

## DSM-IV TR (current standard)

### Autistic Disorder

- A. A total of six (or more) items from (1), (2), and (3), with at least two from (1), and one each from (2) and (3):
- (1) qualitative impairment in social interaction, as manifested by at least two of the following:
    - (a) marked impairment in the use of multiple nonverbal behaviors such as eye-to-eye gaze, facial expression, body postures, and gestures to regulate social interaction
    - (b) failure to develop peer relationships appropriate to developmental level
    - (c) a lack of spontaneous seeking to share enjoyment, interests, or achievements with other people (e.g., by a lack of showing, bringing, or pointing out objects of interest)
    - (d) lack of social or emotional reciprocity
  - (2) qualitative impairments in communication as manifested by at least one of the following:
    - (a) delay in, or total lack of, the development of spoken language (not accompanied by an attempt to compensate through alternative modes of communication such as gesture or mime)
    - (b) in individuals with adequate speech, marked impairment in the ability to initiate or sustain a conversation with others
    - (c) stereotyped and repetitive use of language or idiosyncratic language
    - (d) lack of varied, spontaneous make-believe play or social imitative play appropriate to developmental level
  - (3) restricted repetitive and stereotyped patterns of behavior, interests, and activities, as manifested by at least one of the following:
    - (a) encompassing preoccupation with one or more stereotyped and restricted patterns of interest that is abnormal either in intensity or focus
    - (b) apparently inflexible adherence to specific, nonfunctional routines or rituals
    - (c) stereotyped and repetitive motor mannerisms (e.g., hand or finger flapping or twisting, or complex whole body movements)
    - (d) persistent preoccupation with parts of objects
- B. Delays or abnormal functioning in at least one of the following areas, with onset prior to age 3 years: (1) social interaction, (2) language as used in social communication, or (3) symbolic or imaginative play.
- C. The disturbance is not better accounted for by Rett's Disorder or Childhood Disintegrative Disorder.

### "Pick Six"

Impaired social interaction (two required)

Reciprocity, nonverbal behavior, relationships, shared interests

Plus one of the following communication skill problems required

Eye contact, language, initiate/sustain interactions, imaginative play

(no language impairment required for Asperger's Disorder)

(no cognitive impairment required for Asperger's Disorder)

Plus one of the following repetitive behavior patterns

Routines, stereotypic, restricted interests, sensory, preoccupations

Plus Onset before age 3

And impairment of social **or** language **or** symbolic/imaginative play

**Not Rett's Disorder or Childhood Disintegrative Disorder**

## DSM-5 (May, 2013)

### Autism Spectrum Disorder

Must meet criteria A, B, C, and D:

- A. Persistent deficits in social communication and social interaction across contexts, not accounted for by general developmental delays, and manifest by all 3 of the following:
1. Deficits in social-emotional reciprocity; ranging from abnormal social approach and failure of normal back and forth conversation through reduced sharing of interests, emotions, and affect and response to total lack of initiation of social interaction.
  2. Deficits in nonverbal communicative behaviors used for social interaction; ranging from poorly integrated-verbal and nonverbal communication, through abnormalities in eye contact and body-language, or deficits in understanding and use of nonverbal communication, to total lack of facial expression or gestures.
  3. Deficits in developing and maintaining relationships, appropriate to developmental level (beyond those with caregivers); ranging from difficulties adjusting behavior to suit different social contexts through difficulties in sharing imaginative play and in making friends to an apparent absence of interest in people
- B. Restricted, repetitive patterns of behavior, interests, or activities as manifested by at least two of the following:
1. Stereotyped or repetitive speech, motor movements, or use of objects; (such as simple motor stereotypies, echolalia, repetitive use of objects, or idiosyncratic phrases).
  2. Excessive adherence to routines, ritualized patterns of verbal or nonverbal behavior, or excessive resistance to change; (such as motoric rituals, insistence on same route or food, repetitive questioning or extreme distress at small changes).
  3. Highly restricted, fixated interests that are abnormal in intensity or focus; (such as strong attachment to or preoccupation with unusual objects, excessively circumscribed or perseverative interests).
  4. Hyper- or hypo-reactivity to sensory input or unusual interest in sensory aspects of environment; (such as apparent indifference to pain/heat/cold, adverse response to specific sounds or textures, excessive smelling or touching of objects, fascination with lights or spinning objects).
- C. Symptoms must be present in early childhood (but may not become fully manifest until social demands exceed limited capacities)
- D. Symptoms together limit and impair everyday functioning.

### "All Four Required"

Impaired social and communicative interaction (all three required)

Reciprocity, nonverbal behavior, relationships (shared interests, etc)

Plus any two of the following required

Eye contact, language, initiate/sustain interactions, imaginative play

Body language, understanding, tolerance for change, symbolic play

Ritualized patterns, focus on details

Plus two of the following repetitive behavior patterns

Routines, stereotypic, restricted interests, sensory, preoccupations

Plus Onset in early childhood (not necessarily before age 3) *or later*

**And symptoms together limit and impair everyday functioning**

**Not just General Developmental Delay**